

WHY IS QUANTUM PHYSICS BASED ON COMPLEX NUMBERS?

Felix M. Lev

*Artwork Conversion Software Inc., 1201 Morningside Drive,
Manhattan Beach, CA 90266, USA (E-mail: felixlev@hotmail.com)*

Abstract:

The modern quantum theory is based on the assumption that quantum states are represented by elements of a complex Hilbert space. It is expected that in future quantum theory the number field will be not postulated but derived from more general principles. We consider the choice of the number field in a quantum theory based on a finite field. We assume that the symmetry algebra is the finite field analog of the de Sitter algebra $so(1,4)$ and consider spinless irreducible representations of this algebra. It is shown that the finite field analog of complex numbers is the minimal extension of the residue field modulo p for which the representations are fully decomposable.

1 The statement of the problem

The modern quantum theory is based on the following assumptions:

- *Assumption 1: Quantum states are represented by elements of a (projective) complex Hilbert space.*
- *Assumption 2: Observable physical quantities are represented by selfadjoint operators in this space.*

The field of complex numbers is algebraically closed, i.e. any equation of the n th power in this field has precisely n solutions. As a consequence, any linear operator in a finite dimensional space has at

least one eigenvalue. However, this is not necessarily the case if the space is infinite-dimensional.

The usual motivation of Assumption 2 is that since any physical quantity can take only real values, the spectrum of the corresponding operator should necessarily be real. According to the spectral theorem for selfadjoint operators in Hilbert spaces, this is indeed the case. However, detailed arguments given in Ref. [1] and references therein show that the real spectrum and Assumption 2 are not necessary for constructing meaningful quantum theory. Note that (by definition) any complex number is simply a pair of real numbers, and even for this reason it is not clear why the case of complex spectrum should be excluded. For example, the complex spectrum can represent a pair of real physical quantities.

In quantum theory it is also postulated that the following requirement should be valid:

- *Requirement 1: Any linear operator representing a physical quantity should have a spectral decomposition.*

This implies that one can construct a basis such that any its element is the eigenvector of the given operator. As it is usual in quantum physics, in the general case the basis is understood in the sense of distributions, i.e. points belonging to the continuous spectrum are also treated as eigenvalues.

As follows from the spectral theorem, if one accepts Assumption 2 then Requirement 1 is satisfied automatically. However, the spectral decomposition exists not only for selfadjoint operators; for example, it also exists for unitary operators. It is also clear that the spectral theorem for selfadjoint operators is valid not only in complex Hilbert spaces but in real Hilbert spaces as well. Therefore the only motivation of Assumption 1 is that quantum theory based on complex numbers successfully describes a wide range of physical phenomena.

It is reasonable to believe that in future quantum physics the choice of the number field (or body) will be substantiated instead of saying that a particular number field should be chosen because the

corresponding version of quantum theory is in agreement with experimental data.

In the literature several possibilities have been considered when the principle number field is not the field of complex numbers. There exists a wide literature devoted to quaternionic, p -adic and adelic versions of quantum theory. In each case the theory has its own interesting properties but the problem of the motivation of the choice of the principal number field remains. There also exists a number of works in which implications of finite fields in quantum physics are considered (see e.g. Ref. [2]). However, to the best of our knowledge, only in Refs. [3, 4] and our subsequent publications it has been considered a case when a finite field is the principle number field in quantum theory.

There are several arguments for choosing a finite field as a principal number field in quantum theory is as follows. For example, if one accepts that the ultimate quantum theory should not contain actual infinity at all, then the only possible choice of a number field is the choice of a finite field. It is well known (see e.g. the standard textbooks [5]) that any finite field contains p^n elements, where p is a prime number and n is a natural number. Moreover, the choice of p and n defines the finite field uniquely up to isomorphism. The number p is called the characteristic of the finite field.

We use $GF(p^n)$ to denote a finite field containing p^n elements. As it has been shown in Refs. [3, 4] and our subsequent publications (see e.g. Ref. [6]), a quantum theory based on a Galois field $GF(p^2)$ (GFQT) is a natural generalization of the standard quantum theory based on complex numbers.

Suppose that in our world the principal number field in quantum theory is a finite field characterized by some value of p . Then we still have to answer the question whether there exist deep reasons for choosing this particular value of p or this is simply an accident that our Universe has been created with this value. In any case, if we accept that p is a universal constant then the problem arises what the value of n is. In view of the above discussion it is desirable not to postulate that $n = 2$ but to find a motivation for such a choice. By analogy with

Assumption 1, we accept that

- *Assumption G1: Quantum states in GFQT are represented by elements of a linear projective space over a field $GF(p^n)$ and physical quantities are represented by linear operators in that space.*

Then we do not require any analog of Assumption 2. Instead we accept Requirement 1 which in the case of the GFQT has the same formulation. In the case of finite-dimensional spaces, the existence of the spectral decomposition for some operator A means precisely that one can construct a basis in the usual sense such that all its elements are the eigenvectors of A .

Note that any finite field is not algebraically closed, i.e. any equation of the n th power in the field does necessarily have n solutions. Moreover, it can have no solutions at all. For this reason there is no guarantee that any linear operator in a space over a finite field has even one eigenvalue, to say nothing about the possibility that it has a spectral decomposition.

In the present paper we assume that the symmetry algebra in GFQT is the finite field analog of the de Sitter algebra $so(1,4)$ and consider spinless irreducible representations (IRs) of this algebra. *The main result of the paper is the proof that if $p = 3 \pmod{4}$ then the minimal extension of $GF(p)$ for which there exist 10 linearly independent representation operators satisfying Requirement 1, is the field $GF(p^2)$.*

The paper is organized as follows. In Sect. 2 we describe well known facts about modular representations. We also consider modular IRs of the $su(2)$ algebra and argue that the case $p = 3 \pmod{4}$ is more natural than $p = 1 \pmod{4}$. In Sect. 3 we describe spinless modular IRs of the $so(1,4)$ algebra and in Sect. 4 construct a basis convenient for investigating the spectrum of the operator M_{04} . The main result is proved in Sect. 5, and in Sect. 6 we discuss Hermiticity conditions when IR is supplied by a scalar product. Finally, Sect. 7 is discussion.

2 Modular representations of Lie algebras

In standard quantum theory a Lie algebra over the field of real numbers is called the symmetry algebra if the given system is described by a selfadjoint representation of the algebra in a Hilbert space. For this reason we accept the following

Definition: a representation of the Lie algebra over $GF(p)$ in a space over a finite field is called fully decomposable if the space of representation operators has a basis such that all its elements have a spectral decomposition.

Then a Lie algebra over $GF(p)$ is the symmetry algebra if the system is described by a fully decomposable representation of the algebra in a linear space over a finite field.

As noted in Ref. [4, 6], the correspondence between GFQT and standard quantum theory exists if quantum states in GFQT are represented by elements of a space over $GF(p^2)$ and p is very large. The field $GF(p^2)$ is a quadratic extension of the field $GF(p)$.

It is well known that the element $p - 1 \in GF(p)$, which can be written simply as -1, is a quadratic residue if $p = 1 \pmod{4}$ and quadratic nonresidue if $p = 3 \pmod{4}$. Therefore in the latter case the field $GF(p^2)$ can be treated as a complex extension of $GF(p)$ (i.e. the elements of $GF(p^2)$ can be formally written as $a+bi$ where $a, b \in GF(p)$ and i formally satisfies the condition $i^2 = -1$). The field $GF(p^2)$ has only one nontrivial automorphism which will be denoted as $z \rightarrow \bar{z}$ if $z \in GF(p^2)$. In the case $p = 3 \pmod{4}$, it coincides with the standard complex conjugation $a + bi \rightarrow a - bi$.

By analogy with the conventional quantum theory, one could require that linear spaces V over $GF(p^2)$, used for describing physical states in the GFQT, are supplied by a scalar product $(\dots, \dots)$ such that for any $x, y \in V$ and $a \in GF(p^2)$, (x, y) is an element of $GF(p^2)$ and the following properties are satisfied:

$$(x, y) = \overline{(y, x)}, \quad (ax, y) = \bar{a}(x, y), \quad (x, ay) = a(x, y) \quad (1)$$

In the general case a scalar product in V does not define any positive definite metric, and hence a probabilistic interpretation exists

only for a subset in V [4]. In particular, $(e, e) = 0$ does not necessarily imply that $e = 0$. The quantity (e, e) can be called the norm (or norm squared) of the element e , but in $GF(p)$ the separation of elements into positive and negative does not have the same meaning as in the usual case.

If A_1 and A_2 are linear operators in V such that

$$(A_1x, y) = (x, A_2y) \quad \forall x, y \in V \quad (2)$$

they are said to be adjoint: $A_2 = A_1^*$. Then $A_1^{**} = A_1$ and $A_2^* = A_1$. If $A = A^*$ then, by analogy with the standard cases, we can say that operator A is Hermitian. If $Ae = ae$, $a \in GF(p^2)$ and $e \neq 0$ then the element e is called the eigenvector of the operator A with the eigenvalue a . If $A^* = A$ then by analogy with the standard case $a \in GF(p)$ if $(e, e) \neq 0$. However, if $(e, e) = 0$ then such a conclusion cannot be drawn.

If A is a Hermitian operator such that

$$\begin{aligned} Ae_1 &= a_1e_1, & Ae_2 &= a_2e_2, & (e_1, e_1) &\neq 0, \\ (e_2, e_2) &\neq 0, & a_1 &\neq a_2 \end{aligned} \quad (3)$$

then as in the usual case, $(e_1, e_2) = 0$. We will see below that there also exists a possibility that

$$(e_1, e_1) = (e_2, e_2) = 0 \quad (e_1, e_2) \neq 0 \quad (4)$$

In that case it easy to see from Eqs. (1) and (2) that $\bar{a}_1 = a_2$.

Suppose that the elements $e_1, \dots, e_N$ form a basis in a space over a finite field and there exists a scalar product such that $(e_k, e_l) = 0$ if $k \neq l$ and $(e_k, e_k) \neq 0 \quad \forall k, l = 1, 2, \dots, N$. Then if $x = c_1e_1 + \dots, c_Ne_N$, the coefficient c_k can be found as $(e_k, x)/(e_k, e_k)$.

Representations in spaces over a field of nonzero characteristics are called modular representations. There exists a wide literature devoted to such representations (see e.g. Ref. [7] and references therein). In particular, it has been shown by Zassenhaus [8] that all modular IRs are finite-dimensional and in numerous papers the maximum dimension of such representations is considered.

It is worth noting that usually mathematicians consider only representations over an algebraically closed (infinite) field while our approach is different. We consider only finite fields and investigate what is the minimal extension of $GF(p)$ such that modular IRs of the symmetry algebra are fully decomposable.

Consider, for example, a modular analog of IRs of the $\mathfrak{su}(2)$ algebra. Let $\mathbf{J} = (J_1, J_2, J_3)$ be the operator of ordinary rotations in the standard theory. If $\hbar/2$ rather than $\hbar$ is taken as a unit of measurement of angular momentum then the commutation relations for the components of $\mathbf{J}$ have the form

$$[J_1, J_2] = 2iJ_3, \quad [J_3, J_1] = 2iJ_2, \quad [J_2, J_3] = 2iJ_1 \quad (5)$$

Define the operators $J_{\pm}$ such that

$$J_1 = J_+ + J_- \quad J_2 = -i(J_+ - J_-) \quad (6)$$

Then Eq. (5) can be rewritten as

$$[J_3, J_+] = 2J_+ \quad [J_3, J_-] = -2J_- \quad [J_+, J_-] = J_3 \quad (7)$$

Since Eq. (7) does not contain the quantity i , we now can require that in the modular case the operators $(J_+ J_- J_3)$ act in a space over a finite field and satisfy the same relations.

As follows from Eq. (7), the operator

$$K = J_3^2 - 2J_3 + 4J_+ J_- = J_3^2 + 2J_3 + 4J_- J_+ \quad (8)$$

is the Casimir operator for algebra $(J_+ J_- J_3)$. Consider a representations containing a vector e_0 such that

$$J_+ e_0 = 0, \quad J_3 e_0 = s e_0 \quad (9)$$

where $s \in GF(p)$. Then, as follows from Eq. (8), e_0 is the eigenvector of the operator K with the eigenvalue $s(s+2)$. Denote

$$e_n = (J_-)^n e_0, \quad n = 0, 1, 2, \dots \quad (10)$$

Since K is the Casimir operator, all the e_n are its eigenvectors with the same eigenvalue $s(s+2)$, and, as follows from Eq. (7), $J_3 e_n = (s-2n)e_n$. Hence it follows from Eq. (8) that

$$J_+ J_- e_n = (n+1)(s-n)e_n \quad (11)$$

The maximum value of n , n_{max} is defined by the condition that $J_- e_n = 0$ if $n = n_{max}$. This condition should be compatible with Eq. (11) and therefore $n_{max} = s$. It is easy to see that the elements e_n for $n = 0, 1, \dots, s$ form a basis of modular IR and therefore the dimension of modular IR with a given s is equal to $s+1$, as in the standard case. The only difference is that in the ordinary case s can be any natural number while in the modular case s can take only the values of $0, 1, \dots, p-1$.

In the standard case the operator J_3 is Hermitian while $J_+^* = J_-$. One can assume that the modular IR is considered in a space over $GF(p^2)$ and the same Hermiticity conditions are satisfied. Then it follows from Eq. (11) that

$$(e_{n+1}, e_{n+1}) = (n+1)(s-n)(e_n, e_n) \quad (12)$$

while the elements e_n with the different values of n are orthogonal to each other. Therefore, if $(e_0, e_0) \neq 0$ then all the basis elements have the nonzero norm and are orthogonal to each other. However, we will not assume in advance that modular IRs are considered in a space over $GF(p^2)$. As explained above, our goal is to investigate what is the minimal extension of $GF(p)$ such that modular IRs of the $su(2)$ algebra have three linearly independent observable operators.

The operator J_3 has the spectral decomposition by construction. Consider now the operator $J_1 = J_+ + J_-$ which in the standard theory is the x component of the angular momentum. We use the Pochhammer symbol $(a)_l$ to denote $a(a+1)\cdots(a+l-1)$ and the standard notation

$$F(a, b; c; z) = \sum_l \frac{(a)_l (b)_l z^l}{l! (c)_l} \quad (13)$$

for the hypergeometric series. Let us define

$$e_j^{(x)} = \sum_{k=0}^s \frac{1}{k!} F(-j, -k; -s; 2) e_k \quad (14)$$

As follows from Eqs. (10) and (11),

$$\begin{aligned} J_1 e_j^{(x)} &= \sum_{k=0}^s \frac{1}{k!} \{ F(-j, -k-1; -s; 2)(s-k) + \\ &F(-j, -k+1; -s; 2)k \} e_k \end{aligned} \quad (15)$$

Now we use the following relation between the hypergeometric functions (see e.g. Ref. [9]):

$$\begin{aligned} [c-2a-(b-a)z]F(a, b; c; z) + a(1-z)F(a+1, b; c; z) - \\ (c-a)F(a-1, b; c; z) = 0 \end{aligned} \quad (16)$$

Then it follows from Eq. (15) that $J_1 e_j^{(x)} = (s-2j)e_j^{(x)}$.

A possible way to prove that the elements $e_j^{(x)}$ ($j = 0, 1, \dots, s$) form a basis is to find a transformation inverse to Eq. (14), i.e. to express the elements e_k ($k = 0, 1, \dots, s$) in terms of $e_j^{(x)}$. Let $C_s^j = s!/[j!(s-j)!]$ be the binomial coefficient. Then the transformation has the form

$$e_k = \frac{s!}{2^s(s-k)!} \sum_{j=0}^s C_s^j F(-j, -k; -s; 2) e_j^{(x)} \quad (17)$$

and the proof is as follows. First, as follows from Eq. (14), the r.h.s. of Eq. (17) contains

$$\sum_{j=0}^s C_s^j F(-j, -k; -s; 2) F(-j, -k; -s; 2)$$

We represent this sum as a limit of

$$\sum_{j=0}^s C_s^j F(-j, -k; -s; 2) F(-j, -k; -s; 2) x^j$$

when $x \rightarrow 1$ and use the formula [9]

$$\sum_{j=0}^s C_s^j F(-j, -k; -s; 2) F(-j, -k'; -s; 2) x^j = (1+x)^{s-k-k'} (1-x)^{k+k'} F(-k, -k'; -s; -\frac{4x}{(1-x)^2}) \quad (18)$$

As follows from Eq. (13), the series for the hypergeometric function in Eq. (18) has the last term corresponding to $l = \min(k, k')$ and this term is the most singular when $x \rightarrow 1$. Then it is clear that if $k \neq k'$, the r.h.s. of Eq. (18) is equal to zero in the limit $x \rightarrow 1$ while if $k = k'$ then the limit is equal to $2^s(s-k)!/s!$. This completes the proof of Eq. (17) and we conclude that the operator J_1 has the spectral decomposition even without extending the field $GF(p)$.

Consider now the operator $J_+ - J_-$. Since in the standard theory (see Eq. (6)) it is equal to $-iJ_2$ where J_2 is the y projection of the angular momentum, one might expect that in the modular case $J_+ - J_-$ has a spectral decomposition only if $GF(p)$ is extended.

Consider first the simplest nontrivial case when $s = 1$ ($s = 1/2$ in the standard units). Then, as follows from Eqs. (10) and (11), the characteristic equation for the operator $J_+ - J_-$ is $\lambda^2 = -1$. In the case $p = 3 \pmod{4}$ this equation can be solved only by extending $GF(p)$. However, if $p = 1 \pmod{4}$, the equation has solutions in $GF(p)$ and hence no extension of $GF(p)$ is needed to ensure the spectral decomposition of the operator $J_+ - J_-$. Nevertheless, if p is very large and $p = 1 \pmod{4}$ then the quantities λ satisfying $\lambda^2 = -1 = p - 1$ in $GF(p)$ are very large (at least of order $\sqrt{p}$). This obviously contradicts experimental data since in the IR of the $\mathfrak{su}(2)$ algebra with $s = 1$ no quantities with such large eigenvalues have been observed.

We conclude that the case $p = 1 \pmod{4}$ is probably incompatible with the existing data. For this reason we will consider only quadratic extensions of $GF(p)$ in the case $p = 3 \pmod{4}$. Then by analogy with the above discussion one can prove that the elements

$$e_j^{(y)} = \sum_{k=0}^s \frac{1}{k!} F(-j, -k; -s; 2) i^k e_k \quad (19)$$

are the eigenvectors of the operator $J_+ - J_-$ with the eigenvalues $i(s-2j)$ and they form the basis in the representation space.

Our final conclusions in this section are as follows. If quantum theory is based on a finite field then the number p representing the characteristic of the field is such that $p = 3 \pmod{4}$ rather than $p = 1 \pmod{4}$. Then the complex extension of $GF(p)$ guarantees that modular IRs of the $\mathfrak{su}(2)$ algebra are fully decomposable.

3 Modular IRs of the $\mathfrak{so}(1,4)$ algebra

In standard quantum theory one can choose the units in which $\hbar/2 = c = 1$. Then the assumption that the de Sitter algebra $\mathfrak{so}(1,4)$ is the symmetry algebra implies that its representation operators M^{ab} ($a, b = 0, 1, 2, 3, 4$, $M^{ab} = -M^{ba}$) are Hermitian and satisfy the commutation relations

$$[M^{ab}, M^{cd}] = -2i(\eta^{ac}M^{bd} + \eta^{bd}M^{as} - \eta^{ad}M^{bc} - \eta^{bc}M^{ad}) \quad (20)$$

where η^{ab} is the diagonal metric tensor such that $\eta^{00} = -\eta^{11} = -\eta^{22} = -\eta^{33} = -\eta^{44} = 1$.

One could define the de Sitter invariance in GFQT by saying that the operators M^{ab} describing the system act in a space V over $GF(p^2)$ and satisfy the same relations (20) in that space. However, as noted in Sect. 1, our goal is not to postulate the choice of the number field but substantiate it from the requirement that modular IRs of the symmetry algebra are fully decomposable. Since $\mathfrak{su}(2)$ is a subalgebra of $\mathfrak{so}(1,4)$, it follows from the results of the preceding section, one cannot obtain a fully decomposable IR without extending the field $GF(p)$. However, we do not know yet, whether the complex extension will be sufficient. For this reason it is desirable not to fix the field immediately but assume only that it is an extension of $GF(p)$. In this case we first have to investigate what conclusions can be drawn without assuming the existence of any scalar product and Hermiticity requirements.

In the standard theory, instead of M^{ab} one can work with the set of operators $(\mathbf{J}', \mathbf{J}'', R_{jk})$ ($j, k = 1, 2$). Here $\mathbf{J}'$ and $\mathbf{J}''$ are two

independent $\text{su}(2)$ algebras (i.e. $[\mathbf{J}', \mathbf{J}''] = 0$) described by Eqs. (6) and (7). The commutation relations of the operators $\mathbf{J}'$ and $\mathbf{J}''$ with the R_{jk} have the form

$$\begin{aligned}
[J'_3, R_{1j}] &= R_{1j}, & [J'_3, R_{2j}] &= -R_{2j}, & [J_3'', R_{j1}] &= R_{j1}, \\
[J_3'', R_{j2}] &= -R_{j2}, & [J'_+, R_{2j}] &= R_{1j}, & [J_+'', R_{j2}] &= R_{j1}, \\
[J'_-, R_{1j}] &= R_{2j}, & [J_-'', R_{i1}] &= R_{i2}, & [J'_+, R_{1j}] &= \\
[J_+'', R_{j1}] &= [J'_-, R_{2j}] = [J_-'', R_{j2}] = 0,
\end{aligned} \tag{21}$$

and the commutation relations of the operators R_{jk} with each other have the form

$$\begin{aligned}
[R_{11}, R_{12}] &= 2J'_+, & [R_{11}, R_{21}] &= 2J_+'', \\
[R_{11}, R_{22}] &= -(J'_3 + J_3''), & [R_{12}, R_{21}] &= J'_3 - J_3'', \\
[R_{11}, R_{22}] &= -2J_-'', & [R_{21}, R_{22}] &= -2J'_-
\end{aligned} \tag{22}$$

Then, if $\mathbf{M} = \{M^{23}, M^{31}, M^{12}\}$, $\mathbf{B} = -\{M^{14}, M^{24}, M^{34}\}$ and the relation between the sets $(\mathbf{J}', \mathbf{J}'', R_{jk})$ and M^{ab} is given by

$$\begin{aligned}
\mathbf{M} &= \mathbf{J}' + \mathbf{J}'', & \mathbf{B} &= \mathbf{J}' - \mathbf{J}'', & M_{01} &= i(R_{11} - R_{22}), \\
M_{02} &= R_{11} + R_{22}, & M_{03} &= -i(R_{12} + R_{21}), \\
M_{04} &= R_{12} - R_{21},
\end{aligned} \tag{23}$$

one can directly verify that Eq. (20) follows from Eqs. (7), (21), (22), (23) and *vice versa*.

Since Eqs. (7), (21) and (22) do not contain the quantity i , one can define the de Sitter invariance in the modular case by requiring that the system is described by the operators $(\mathbf{J}', \mathbf{J}'', R_{jk})$ ($j, k = 1, 2$) satisfying these expressions. At first glance, these relations might seem rather chaotic, but they are natural in the Weyl basis of the $\text{so}(1,4)$ algebra.

Proceeding from the method of $\text{su}(2) \times \text{su}(2)$ shift operators, developed by Hughes [10] for constructing standard unitary IRs of the group $\text{SO}(5)$, and following Ref. [4], we now give a description of modular IRs of the $\text{so}(1,4)$ algebra.

Consider the space of maximal $su(2) \times su(2)$ vectors, i.e. such vectors x that $J'_+ x = J_+'' x = 0$. Then it follows from Eqs. (7), (21), (22) that the operators

$$\begin{aligned} A^{++} &= R_{11} \quad A^{+-} = R_{12}(J_3'' + 1) - J_-'' R_{11}, \\ A^{-+} &= R_{21}(J'_3 + 1) - J'_- R_{11} \\ A^{--} &= -R_{22}(J'_3 + 1)(J_3'' + 1) + J_-'' R_{21}(J'_3 + 1) + \\ &J'_- R_{12}(J_3'' + 1) - J'_- J_-'' R_{11} \end{aligned} \quad (24)$$

act invariantly on this space. The notations are related to the property that if x^{kl} ($k, l > 0$) is the maximal $su(2) \times su(2)$ vector and simultaneously the eigenvector of operators J'_3 and J_3'' with the eigenvalues k and l , respectively, then $A^{++}x^{kl}$ is the eigenvector of the same operators with the values $k + 1$ and $l + 1$, $A^{+-}x^{kl}$ - the eigenvector with the values $k + 1$ and $l - 1$, $A^{-+}x^{kl}$ - the eigenvector with the values $k - 1$ and $l + 1$ and $A^{--}x^{kl}$ - the eigenvector with the values $k - 1$ and $l - 1$.

As follows from the results of the preceding section, the vector $x_{\alpha\beta}^{kl} = (J'_-)^{\alpha}(J_-'')^{\beta}x^{kl}$ is the eigenvector of the operators J'_3 and J_3'' with the eigenvalues $k - 2\alpha$ and $l - 2\beta$, respectively. Since

$$\mathbf{J}^2 = J_3'^2 - 2J_3 + 4J_+J_- = J_3''^2 + 2J_3 + 4J_-J_+$$

is the Casimir operator for the $\mathbf{J}$ algebra, it follows in addition that

$$\mathbf{J}'^2 x_{\alpha\beta}^{kl} = k(k + 2)x_{\alpha\beta}^{kl}, \quad \mathbf{J}''^2 x_{\alpha\beta}^{kl} = l(l + 2)x_{\alpha\beta}^{kl} \quad (25)$$

$$J'_+ x_{\alpha\beta}^{kl} = \alpha(k + 1 - \alpha)x_{\alpha-1,\beta}^{kl}, \quad J_+'' x_{\alpha\beta}^{kl} = \beta(l + 1 - \beta)x_{\alpha,\beta-1}^{kl} \quad (26)$$

From these formulas it follows that the action of the operators $\mathbf{J}'$ and $\mathbf{J}''$ on x^{kl} generates a space with the dimension $(k + 1)(l + 1)$ and the basis $x_{\alpha\beta}^{kl}$ ($\alpha = 0, 1, \dots, k$, $\beta = 0, 1, \dots, l$).

The Casimir operator of the second order for the algebra (20) can be written as

$$\begin{aligned} I_2 &= -\frac{1}{2} \sum_{ab} M_{ab} M^{ab} = \\ &4(R_{22}R_{11} - R_{21}R_{12} - J'_3) - 2(\mathbf{J}'^2 + \mathbf{J}''^2) \end{aligned} \quad (27)$$

and the basis in the representation space can be explicitly constructed assuming that there exists a vector e^0 which is the maximal $\text{su}(2) \times \text{su}(2)$ vector such that

$$\begin{aligned} \mathbf{J}'e^0 &= J_+''e^0 = 0, \quad J_3''e^0 = se^0, \\ I_2e^0 &= [w - s(s+2) + 9]e^0 \end{aligned} \quad (28)$$

(see Ref. [4] for details) where $w, s \in GF(p)$.

Define the vectors

$$e^{nr} = (A^{++})^n (A^{+-})^r e^0 \quad (29)$$

Then a direct calculation using Eqs. (7), (21), (22), (24), (27), (28), and (29) gives

$$A^{--}A^{++}e^{nr} = -\frac{1}{4}(n+1)(n+s+2)[w + (2n+s+3)^2]e^{nr} \quad (30)$$

$$A^{-+}A^{+-}e^{nr} = -\frac{1}{4}(r+1)(s-r)[w + 1 + (2r-s)(2r+2-s)]e^{nr} \quad (31)$$

As follows from the last expression, r can take only the values of $0, 1, \dots, s$, as well as in the standard theory. At the same time, as follows from Eq. (30), the possible values of n in the modular case are not the same as in the standard theory. Indeed, in the standard case the possible values of n are obviously $n = 0, 1, 2, \dots, \infty$ and therefore IR is infinite dimensional. On the contrary, in the modular case n can take only the values $0, 1, \dots, n_{max}$ where the maximum value of n , n_{max} can be found as follows. By definition of the operator A^{++} , $A^{++}e^{nr} = 0$ if $n = n_{max}$. This relation should not contradict Eq. (30) if $n = n_{max}$. Therefore n_{max} is the least number satisfying the congruence modulo p

$$(n_{max} + 1)(n_{max} + s + 2)[w + (2n_{max} + s + 3)^2] = 0 \quad (32)$$

In particular, the IR is necessarily finite dimensional in agreement with the Zassenhaus theorem [8].

In the standard theory $w = \mu^2$ where μ is the particle dS mass (see e.g. Ref. [11]) but in GFQT the element $w \in GF(p)$ is not

necessarily a square in $GF(p)$. Since -1 is a quadratic residue in $GF(p)$ if $p \equiv 1 \pmod{4}$ and quadratic nonresidue if $p \equiv 3 \pmod{4}$ then in the latter case (which is of only interest for us in view of the results of the preceding section) the equality $w + (2n_{max} + s + 3)^2 = 0$ in $GF(p)$ is possible only if w is a quadratic nonresidue. We will see below that only this condition is consistent. Then $w = -\tilde{\mu}^2$ where for $\tilde{\mu}$ obviously two solutions are possible.

Consider for simplicity the case $s = 0$. Then n_{max} should satisfy one of the conditions $2n_{max} + 3 = \pm\tilde{\mu}$ and therefore there exist two solutions for n_{max} . We should choose one with the lesser value of n_{max} and, as follows from Eq. (32), this value should be less than $p - 2$. Let us assume that both, $\tilde{\mu}$ and $-\tilde{\mu}$ are represented by $0, 1, \dots, (p - 1)$. Then if $\tilde{\mu}$ is odd, $-\tilde{\mu} = p - \tilde{\mu}$ is even and *vice versa*. We choose the odd number as $\tilde{\mu}$. Then the two solutions are $n_1 = (\tilde{\mu} - 3)/2$ and $n_2 = p - (\tilde{\mu} + 3)/2$. It is obvious that $n_1 < n_2$ and $n_1 < p - 2$. Therefore

$$n_{max} = (\tilde{\mu} - 3)/2 \quad (33)$$

In particular, this quantity satisfies the condition $n_{max} \leq (p - 5)/2$.

Since e^{nr} is the maximal $su(2) \times su(2)$ vector with the eigenvalues of the operators $\mathbf{J}'$ and $\mathbf{J}''$ equal to $n + r$ and $n + s - r$, respectively, then as a basis of the representation space one can take the vectors $e_{\alpha\beta}^{nr} = (J'_-)^{\alpha}(J_-'')^{\beta}e^{nr}$ where, for the given n and s , the quantity α can take the values of $0, 1, \dots, n + r$ and β - the values of $0, 1, \dots, n + s - r$.

If $s = 0$ then there exist only the maximal $su(2) \times su(2)$ vectors x^{kl} with $k = l$ and therefore the basis of the representation space is formed by the vectors $e_{\alpha\beta}^n \equiv e_{\alpha\beta}^{n0}$ where $n = 0, 1, 2, \dots$; $\alpha, \beta = 0, 1, \dots, n$. The explicit expressions for the action of the operators R_{jk} in this basis

can be calculated by using Eqs. (21), (22) (29-31) and the result is

$$\begin{aligned}
R_{11}e_{\alpha\beta}^n &= \frac{(n+1-\alpha)(n+1-\beta)}{(n+1)^2}e_{\alpha\beta}^{n+1} + \\
&\frac{\alpha\beta n}{4(n+1)}[w+(2n+1)^2]e_{\alpha-1,\beta-1}^{n-1}, \\
R_{12}e_{\alpha\beta}^n &= \frac{n+1-\alpha}{(n+1)^2}e_{\alpha,\beta+1}^{n+1} - \frac{\alpha n}{4(n+1)}[w+(2n+1)^2]e_{\alpha-1,\beta}^{n-1}, \\
R_{21}e_{\alpha\beta}^n &= \frac{n+1-\beta}{(n+1)^2}e_{\alpha+1,\beta}^{n+1} - \frac{\beta n}{4(n+1)}[w+(2n+1)^2]e_{\alpha,\beta-1}^{n-1}, \\
R_{22}e_{\alpha\beta}^n &= \frac{1}{(n+1)^2}e_{\alpha+1,\beta+1}^{n+1} + \\
&\frac{n}{4(n+1)}[w+(2n+1)^2]e_{\alpha\beta}^{n-1}
\end{aligned} \tag{34}$$

We can now prove that the only consistent case when Eq. (32) is satisfied is that w is a quadratic nonresidue and $n_{max} = p - 2$ is impossible. Indeed, if n_{max} is a maximum possible value of n then, as follows from Eq. (34), $R_{11}e^{n_{max}} = R_{12}e^{n_{max}} = 0$. Since $e^{n_{max}} = e_{00}^{n_{max}}$, it follows from Eq. (27), $I_2e^{n_{max}} = -4n_{max}(n_{max}+3)e^{n_{max}}$. This condition can be compatible with $I_2e^{n_{max}} = (w+9)e^{n_{max}}$ (see Eq. (28)) only if $w + (2n_{max} + 3)^2 = 0$.

As follows from the results of the preceding section, from $\mathbf{J}'$ and $\mathbf{J}''$ one can construct six linearly independent operators having a spectral decomposition if $GF(p)$ is extended to $GF(p^2)$ in the case $p = 3 \pmod{4}$. Our goal is to prove that such an extension is sufficient to ensure a possibility of constructing additional four independent operators from the R_{jk} such that all of them have a spectral decomposition.

For simplicity we consider only the spinless case. Then, as follows from Eq. (28), the vector e_0 is annihilated by all the representation operators of the $so(4)=su(2)\times su(2)$ algebra. Therefore all the operators $M_{0\nu}$ ($\nu = 1, 2, 3, 4$) are on equal footing and it is sufficient to prove that the operator M_{04} has a spectral decomposition.

4 M_{04} operator in the $(\mathbf{J}^2, J_3)$ basis

We now use $\mathbf{J}$ to denote $\mathbf{J}' + \mathbf{J}''$. In the standard theory $\mathbf{J}$ is the angular momentum corresponding to conventional three-dimensional rotations. In the modular case the set $\mathbf{J}$ is defined by the operators $(J_+ J_- J_3)$ satisfying the commutation relations (7) since $\mathbf{J}'$ and $\mathbf{J}''$ satisfy these relations and $[\mathbf{J}', \mathbf{J}''] = 0$. Since $M_{04} = R_{12} - R_{21}$ (see Eq. (23)) then, as follows from Eq. (21), $[M_{04}, \mathbf{J}] = 0$. Therefore for investigating operator M_{04} it is convenient to decompose the representation space into subspaces such that all the elements belonging to the same subspace are the eigenvectors of the operators $\mathbf{J}^2$ and J_3 with the same eigenvalue.

Since e^n satisfies the conditions $J'_+ e^n = J_+'' e^n = 0$ by construction, it also satisfies $J_+ e^n = 0$ and $J_3 e^n = 2n e^n$. Therefore the elements $(J_-)^l e^n$ ($l = 0, 1, \dots, 2n$) form a subspace corresponding to IR of the $\text{su}(2)$ algebra with the spin $s = 2n$.

Let now $e(n, k)$ be an element satisfying the conditions $J_+ e(n, k) = 0$ and $J_3 e(n, k) = 2(n - k) e(n, k)$. Then the elements $e(n, k, l) = (J_-)^l e(n, k)$ ($l = 0, 1, \dots, 2(n - k)$) form a subspace corresponding to IR of the $\text{su}(2)$ algebra with the spin $s = 2(n - k)$. We define

$$e(n, k) = B(n, k) e(n, 0) \quad (35)$$

where $e(n, 0) = e^n$ and

$$B(n, k) = \frac{1}{(n!)^2} \sum_{l=0}^k (-1)^{k-l} C_k^l (n-l)! (n+l-k)! (J_-')^l (J_-'')^{k-l} \quad (36)$$

Then we have to prove that $J_+ e(n, k) = 0$ for $k = 1, 2, \dots, n$. A direct calculation using Eqs. (7), (35) and (36) gives

$$J'_+ e(n, k) = k e(n, k-1) \quad J_+'' e(n, k) = -k e(n, k-1) \quad (37)$$

Therefore

$$(J'_+ + J_+'') e(n, k) = 0 \quad (J'_- - J_+'') e(n, k) = 2k e(n, k-1) \quad (38)$$

It has been shown in the preceding section, that acting by J'_- and J_+'' on the element e^n one can obtain a subspace with the

dimension $(n+1)^2$ and the basis $e_{\alpha\beta}^n$. On the other hand, as shown in this section, in such a way it is also possible to obtain a subspace with the basis $e(n, k, l)$ where at fixed n and k , l takes the values $0, 1, \dots, 2(n-k)$ and at a fixed n , k takes the values $0, 1, \dots, n$. The problem arises whether the elements $e(n, k, l)$ ($n = 0, 1, \dots, n_{max}$) also form a basis in the representation space.

In the standard theory the proof follows from the fact that the dimension of the subspace generated by the elements $e(n, k, l)$ at different values of k and l is also equal to $(n+1)^2$ since

$$\sum_{k=0}^n [2(n-k) + 1] = (n+1)^2$$

Moreover, as follows from Eq. (38), any element $e(n, k, l)$ can be chosen as a cyclic element of IR. It will also be shown in Sect. 6 that it is possible to define a scalar product in the representation space such that the both basis's, $\{e_{\alpha\beta}^n\}$, and $\{e(n, k, l)\}$ satisfy the following property: their different elements are orthogonal while the norm of each element is not equal to zero.

In the modular case the situation might be a bit more complicated. As shown in Sect. 2, the dimension of modular IR of the $su(2)$ algebra characterized by the value J is equal to $J+1$ only if J is one of the values $0, 1, \dots, p-1$. Therefore the dimension of IR characterized by the values of n and k is $2(n-k) + 1$ only if $(n-k) \leq (p-1)/2$. For example, if $n-k = (p+1)/2$, IR corresponds to $J = 1$ and has the dimension 2. This examples shows that in the modular case J is not necessarily even in the spinless case.

As noted in the preceding section, the quantity w should be a quadratic nonresidue and then the value of n_{max} is necessarily less than $(p-3)/2$. In that case all the possible values of J are even and the dimension of IR characterized by the maximal weight J is $J+1$, as well as in the standard theory. Therefore the subspace generated by the elements $e(n, k, l)$ at different values of k and l also has the dimension $(n+1)^2$ and all the the elements $e(n, k, l)$ form a basis in the representation space.

Let $V(j)$ be a subspace generated by the elements $e(n, k)$ such that the value of $j = J/2 = (n - k)$ is the same for all the basis elements. Then they satisfy the conditions $J_+e(n, k) = 0$ and $J_3e(n, k) = 2je(n, k)$. The basis in $V(j)$ is formed by the elements $e(n, n - j)$ ($n = j, j + 1, \dots, n_{max}$). One can also define subspaces $V(j, k) = (J_-)^k V(j)$. It is clear that the representation space can be decomposed into the subspaces $V(j, k)$ such that $k = 0, 1, \dots, 2j$ if j is fixed and $j = 0, 1, \dots, n_{max}$. Since M_{04} commutes with $\mathbf{J}$, each subspace $V(j, k)$ is invariant under the action of M_{04} and for a fixed j the spectrum of M_{04} in all the subspaces $V(j, k)$ is the same. Therefore for investigating operator M_{04} it is sufficient to consider its action in subspaces $V(j)$.

Suppose that j is fixed and denote $E_n = (-1)^n e(n + j, n)$. Then the elements E_n ($n = 0, 1, \dots, n_{max} - j$) form a basis in $V(j)$. A direct calculation using Eqs. (22) and (34-36) shows that the action of $M_{04} = A$ in $V(j)$ is given by

$$AE_n = E_{n+1} + \frac{n(n + 2j + 1)}{4(n + j)(n + j + 1)}[w + (2n + 2j + 1)^2]E_{n-1} \quad (39)$$

This expression shows that the matrix of the operator A has only the following nonzero elements:

$$A_{n+1, n} = 1 \quad A_{n, n+1} = \frac{(n + 1)(n + 2j + 2)}{4(n + j + 1)(n + j + 2)}[w + (2n + 2j + 3)^2] \quad (40)$$

where $n = 0, 1, \dots, n_{max} - j$.

Note that the results of this and preceding sections have been obtained assuming that IR is considered in a space over a finite field of characteristic p but no concrete choice of the field with such a characteristic has been made.

5 Spectrum of the M_{04} operator

Consider now the spectrum of the operator having the matrix (40). Let $A(\lambda)$ be the matrix of the operator $A - \lambda$. We use $\Delta_k^l(\lambda)$ to denote the

determinant of the matrix obtained from $A(\lambda)$ by taking into account only the rows and columns with the numbers $k, k+1, \dots, l$. Our convention is that in the matrix $A(\lambda)$ the first row and column have the values equal to 0 while the last ones have the values equal to $N = n(j)_{max}$ which should be defined yet. Therefore the characteristic equation can be written as

$$\Delta_0^N(\lambda) = 0 \quad (41)$$

The matrix $A(\lambda)$ is three-diagonal. It is easy to see that

$$\Delta_0^{n+1}(\lambda) = -\lambda \Delta_0^n(\lambda) - A_{n,n+1} A_{n+1,n} \Delta_0^{n-1}(\lambda) \quad (42)$$

Let λ_l be a solution of Eq. (41). Then the element

$$\chi(\lambda_l) = \sum_{n=0}^N \{(-1)^n \Delta_0^{n-1}(\lambda_l) E_n / [\prod_{k=0}^{n-1} A_{k,k+1}]\} \quad (43)$$

is the eigenvector of the operator A with the eigenvalue λ_l . This can be verified directly by using Eqs. (39-43).

To solve Eq. (41) we have to find the expressions for $\Delta_0^n(\lambda)$ when $n = 0, 1, \dots, N$. It is obvious that $\Delta_0^0(\lambda) = -\lambda$, and as follows from Eq. (40),

$$\Delta_0^1(\lambda) = \lambda^2 - \frac{w + (2j+3)^2}{2(j+2)} \quad (44)$$

Since w should be a quadratic nonresidue, it can be represented as $w = -\tilde{\mu}^2$. Then it can be shown that $\Delta_0^n(\lambda)$ is given by the following expressions. If n is odd then

$$\begin{aligned} \Delta_0^n(\lambda) = & \sum_{l=0}^{(n+1)/2} C_{(n+1)/2}^l \prod_{k=1}^l [\lambda^2 + (\tilde{\mu} - 2j - 4k + 1)^2] (-1)^{(n+1)/2-l} \\ & \prod_{k=l+1}^{(n+1)/2} \frac{2j + 2k + 1}{2(j + (n+1)/2 + k)} (\tilde{\mu} - 2j - 4k + 1)(\tilde{\mu} - 2j - 4k - 1) \end{aligned} \quad (45)$$

and if n is even then

$$\Delta_0^n(\lambda) = (-\lambda) \sum_{l=0}^{n/2} C_{n/2}^l \prod_{k=1}^l [\lambda^2 + (\tilde{\mu} - 2j - 4k + 1)^2] (-1)^{n/2-l} \prod_{k=l+1}^{(n+1)/2} \frac{2j + 2k + 1}{2(j + n/2 + k + 1)} (\tilde{\mu} - 2j - 4k - 1)(\tilde{\mu} - 2j - 4k - 3) \quad (46)$$

Indeed, for $n = 0$ Eq. (46) is compatible with $\Delta_0^0(\lambda) = -\lambda$, and for $n = 1$ Eq. (45) is compatible with Eq. (44). Then one can directly verify that Eqs. (45) and (46) are compatible with Eq. (42).

As noted in the preceding section, N should be such that $N \leq n_{max} - j$ where $w + (2n_{max} + 3)^2 = 0$ in the spinless case. On the other hand, N is the greatest value of n for which the coefficient in front of E_{n-1} in Eq. (39) is not equal to zero. Therefore

$$(N + 2j + 2)[w + (2N + 2j + 3)^2] = 0 \quad (47)$$

(compare with Eq. (33)). As a consequence of the definition of n_{max} , the second multiplier in this expression is equal to zero if $N = n_{max} - j$. Therefore the quantity N is the lesser of $n_{max} - j$ and $p - 2j - 2$. Since $j \leq n_{max}$, the only possibility for N is such that

$$\tilde{\mu} = 2N + 2j + 3 \quad (48)$$

Then, as follows from Eqs. (45) and (46), when N is odd or even, only the term with $l = [(N + 1)/2]$ (where $[(N + 1)/2]$ is the integer part of $(N + 1)/2$) contributes to the sum.

As a consequence

$$\Delta_0^N(\lambda) = (-\lambda)^{r(N)} \prod_{k=1}^{[(N+1)/2]} [\lambda^2 + (\tilde{\mu} - 2j - 4k + 1)^2] \quad (49)$$

where $r(N) = 0$ if N is odd and $r(N) = 1$ if N is even. If $p = 3 \pmod{4}$, this equation has solutions only if $GF(p)$ is extended, and the minimum extension is $GF(p^2)$. Then the solutions are given by

$$\lambda = \pm i(\tilde{\mu} - 2j - 4k + 1) \quad (k = 1, 2, \dots, [(N + 1)/2]) \quad (50)$$

and when N is even there also exists an additional solution $\lambda = 0$. When N is odd (and the dimension of $V(j)$ is even) the solutions can be represented as

$$\lambda = \pm 2i, \pm 6i, \dots \pm 2iN \quad (51)$$

while when N is even, the solutions can be represented as

$$\lambda = 0, \pm 4i, \pm 8i, \dots \pm 2iN \quad (52)$$

Therefore the spectrum is equidistant and the distance between the neighboring elements is equal to $4i$. As follows from Eqs. (48), all the roots are simple. Then, as follows from Eq. (43), the operator M_{04} has a spectral decomposition and this completes the proof of our main statement (see Sect. 1).

6 Hermiticity conditions

As shown in the preceding sections, for physically meaningful modular IRs of the $so(1,4)$ algebra, the extension of $GF(p)$ to $GF(p^2)$ guarantees that they are fully decomposable. Therefore, as explained in Sect. 2, one can define a scalar product in the representation space. By analogy with the standard theory, we now assume that the operators M^{ab} are Hermitian with respect to the chosen scalar product. Therefore, as follows from Eq. (22), the Hermiticity conditions for the operators $(\mathbf{J}', \mathbf{J}'', R_{kl})$ are as follows

$$(J'_-)^* = J'_+ \quad (J_-'')^* = J_+'' \quad R_{12}^* = R_{21} \quad R_{11}^* = -R_{22} \quad (53)$$

while the operators J'_3 and J_3'' should be Hermitian.

In the spinless case, as follows from Eqs. (24), (27), (28) and (30),

$$(e^{n+1}, e^{n+1}) = \frac{n+1}{4(n+2)}[w + (2n+3)^2](e^n, e^n) \quad (54)$$

Therefore, if we assume that $(e^0, e^0) = 1$ then

$$(e^n, e^n) = \frac{1}{4^n(n+1)} \prod_{l=1}^n [w + (2l+3)^2] \quad (55)$$

It is also easy to show that the elements e^n with the different values of n are mutually orthogonal.

Our conclusion is as follows. The scalar products of the basis elements are fully defined by the value of (e^0, e^0) assuming that the operators M^{ab} are Hermitian. If $(e^0, e^0) \neq 0$ then all the basis elements $e_{\alpha\beta}^n$ have a nonzero norm and are mutually orthogonal. This is in agreement with the properties of the scalar product in the modular case (see Sect. 2). The quantum numbers $(n\alpha\beta)$ characterize the eigenvalues of the operators J'_3 , J_3'' and $\mathbf{J}'^2$ (note that in the spinless case the elements $e_{\alpha\beta}^n$ are the eigenvectors of the operators $\mathbf{J}'^2$ and $\mathbf{J}''^2$ with the same eigenvalues).

Our next goal is to show that the elements $e(n, k, l)$ (see Sect. 4) have nonzero norms and are mutually orthogonal. A direct calculation using Eqs. (7), (11), (35), (36) and (55) gives

$$(e(n, k), e(n, k)) = k! \left[\frac{(n-k)!}{n!} \right]^2 \frac{(2n+1-k)!}{(2n+1-2k)!} (e^n, e^n) \quad (56)$$

Then, as follows from the definition of the elements E_n and Eq. (40),

$$(E_n, E_n) = \frac{1}{4^j(j+1)} \left\{ \prod_{l=1}^j [w + (2l+1)^2] \right\} \prod_{l=0}^{n-1} A_{l,l+1} \quad (57)$$

In particular, these elements have nonzero norms and are mutually orthogonal.

Suppose now that λ is one of the eigenvalues given by Eq. (50) and $\chi(\lambda)$ is the eigenvector of M_{04} with this eigenvalue (see Eq. (43)). Since M_{04} is Hermitian and λ is imaginary, the only possible value of $(\chi(\lambda), \chi(\lambda))$ is zero (see Sect. 2). Moreover, since all the eigenvalues are imaginary when N is odd and there also exists an additional eigenvalue $\lambda = 0$ when N is even, $(\chi(\lambda_1), \chi(\lambda_2))$ is necessarily equal to zero if $\lambda_1 + \lambda_2 = 0$. Let us show, however, that $(\chi(\bar{\lambda}), \chi(\lambda)) \neq 0$ for imaginary eigenvalues and $(\chi(0), \chi(0)) \neq 0$.

As follows from Eqs. (43) and (55),

$$(\chi(\bar{\lambda}), \chi(\lambda)) = \frac{1}{4^j(j+1)} \left\{ \prod_{l=1}^j [w + (2l+1)^2] \right\} \left\{ \sum_{n=0}^n \Delta_0^{n-1}(\lambda)^2 / \left[\prod_{k=0}^{n-1} A_{k,k+1} \right] \right\} \quad (58)$$

and one can directly verify that a generalization of Eq. (42) is

$$\Delta_0^N(\lambda) = \Delta_0^n(\lambda) \Delta_{n+1}^N(\lambda) - A_{n,n+1} \Delta_0^{n-1}(\lambda) \Delta_{n+2}^N(\lambda) \quad (59)$$

since in our case $A_{n+1,n} = 1$. Since λ is the eigenvalue, $\Delta_0^N(\lambda) = 0$ and one can use Eq. (59) for $k = n, n-1, \dots, 0$. As a consequence,

$$\Delta_0^n(\lambda) = \left\{ \prod_{l=0}^n A_{l,l+1} \right\} \Delta_{n+2}^N(\lambda) / \Delta_1^N(\lambda) \quad (60)$$

and Eq. (58) can be rewritten as

$$(\chi(\bar{\lambda}), \chi(\lambda)) = \frac{1}{4^j(j+1)} \left\{ \prod_{l=1}^j [w + (2l+1)^2] \right\} \left\{ \sum_{n=0}^n \Delta_0^{n-1}(\lambda) \Delta_{n+1}^N(\lambda) \right\} / \Delta_1^N(\lambda) \quad (61)$$

The sum can be written as $-d\Delta_0^N(\lambda)/d\lambda$ and therefore, as follows from Eqs. (49) and (61)

$$(\chi(\bar{\lambda}), \chi(\lambda)) = \frac{(-\lambda)^{r(N)}}{4^j(j+1)} \left\{ \prod_{l=1}^j [w + (2l+1)^2] \right\} \left\{ \prod_{\lambda_l \neq \lambda} (\lambda - \lambda_l) \right\} / \Delta_1^N(\lambda) \quad (62)$$

where the last product is taken over all the eigenvalues excepting λ . Since all the eigenvalues are simple (see Sect. 5) this product is not equal to zero, and since the l.h.s. of Eq. (62) cannot be singular by construction, it is not equal to zero.

We conclude that the basis elements $\chi(\lambda_l)$ in $V(j)$ satisfy the following orthogonality properties. If λ_l is imaginary then $\bar{\lambda}_l$ also is the eigenvalue. The element $\chi(\lambda_l)$ is orthogonal to itself and all the other elements $\chi(\lambda_k)$ if $\lambda_l \neq \bar{\lambda}_k$ while $(\chi(\lambda_l), \chi(\bar{\lambda}_l)) \neq 0$. When N is even, there also exists the element $\chi(0)$ which is orthogonal to all the other elements $\chi(\lambda_l)$ while $(\chi(0), \chi(0)) \neq 0$.

7 Conclusion

The main difference between our approach and the standard one is that we do not postulate from the beginning that physical states are elements of a specific linear space. Following our previous publications, we assume that the ultimate quantum theory will be based on a finite field. Then we investigate what is the minimum extension of the residue field modulo p such that representations of the symmetry algebra are fully decomposable.

When the characteristic of the finite field p is large, the operators representing physical quantities act in spaces, the dimensions of which are large. One can show [3, 4] that the dimensions of spinless IRs are of order p^3 . In the present paper we decompose the representation space into subspaces $V(j, k)$ the dimensions of which may be of order p . Since finite fields are not algebraically closed, there is no guarantee that the characteristic equation of such a large power will have solutions in the given finite field.

We believe, it is a very interesting result that if the symmetry algebra is the modular analog of the de Sitter algebra $so(1,4)$ then the complex extension is already sufficient for ensuring spinless IRs to be fully decomposable. This might also be an explanation of the fact that the present quantum theory is based on complex numbers.

In the literature the operator M_{04} is usually treated as the de Sitter analog of the energy operator (see e.g. Refs. [11]). For this reason one might think that the existence of imaginary eigenvalues of this operator excludes a possibility that a theory based on a finite field might be realistic. This problem will be discussed elsewhere.

In the standard theory the role of the scalar product is at least threefold: i) to ensure real eigenvalues of selfadjoint operators; ii) to ensure spectral decomposition for such operators; iii) to ensure probabilistic interpretation in Copenhagen formulation. As it has been already noted, there are no reasons of why complex eigenvalues should be excluded, and it has also been shown that ii) can be valid without assuming the existence of any scalar product and Hermiticity requirement. The results of Sect. 6 show that in GFQT one can define a scalar product in such a way that at least for a subset of elements from the representation space, the probabilistic interpretation is valid. However, the problem arises how to interpret the states which do not have counterparts in the standard theory (e.g. the states with zero norm).

Acknowledgements: The author is grateful to H. Doughty, B. Fridman, C. Hayzelden, V.A. Karmanov, G. Mullen, and M.A. Olshanetsky for valuable discussions.

References

- [1] J-M. Levy-Leblond, Who is afraid of non-Hermitian operators? Phase and angle in quantum mechanics, Ann. Phys. (N.Y.) 101 (1976) 319-341; C.M. Bender, D.C. Brody and H.F. Jones, Must a Hamiltonian be Hermitian?, Am. J. Phys. 71 (2003) 1095-1102; A. Mostafazadeh, Pseudo-Hermiticity and generalized PT and CPT symmetries, J. Math. Phys. 44 (2003) 974-989.
- [2] E. Beltrametti, Rotations and Lorentz group in a finite geometry, J. Math. Phys. 9 (1968) 1027-1035; E. Beltrametti, Dirac spinors, covariant currents and the Lorentz group over a finite geometry, Nuovo Cimento 55A (1968) 301-317; H. Doughty, Private Communication of June 24th, 2004; A. Vourdas, Quantum systems with finite Hilbert space, Rep. Progr. Phys. 67 (2004) 267-320; M. Planat and M. Saniga, Abstract algebra, projective geometry and time encoding of quantum information, quant-ph/0503159; M. Planat and

- H. Rosu, Mutually unbiased phase states, phase uncertainties and Gauss sums, Eur. Phys. J., Vol. D, to be published.
- [3] F.M. Lev, Representations of de Sitter algebras over a finite field and their possible interpretations, Sov. J. Nucl. Phys. 48 (1988) 903-912; J. Math. Phys., Modular representations as a possible basis of finite physics, 30 (1989) 1985-1998.
 - [4] F.M. Lev, Finiteness of physics and its possible consequences, J. Math. Phys. 34 (1993) 490-527.
 - [5] B.L. Van der Waerden, Algebra I, Springer-Verlag, Berlin - Heidelberg - New York, 1967; K. Ireland and M. Rosen, A Classical Introduction to Modern Number Theory, Graduate Texts in Mathematics-87, New York - Heidelberg - Berlin, Springer, 1987.
 - [6] F.M. Lev, Massless elementary particles in a quantum theory over a Galois field, Theor. Math. Phys. 138 (2004) 208-225.
 - [7] J.C. Jantzen, Representation of Lie algebras in prime characteristics, NATO Asi series, series C, Math. Phys. Sci. 514 "Representation theories and algebraic geometry", 185-235, Kluwer Acad. Publ. Dordrecht, 1998.
 - [8] H. Zassenhaus, The representations of Lie algebras of prime characteristics, Proc. Glasgow Math. Assoc. 2 (1954) 1-36.
 - [9] H. Bateman and A. Erdelyi, Higher Transcendental Functions, McGraw-Hill Book Company, New York, 1953.
 - [10] W.B. Hughes, $SU(2)*SU(2)$ shift operators and representations of $SO(5)$, J. Math. Phys. 24 (1983) 1015-1020.
 - [11] F.M. Lev, Could only fermions be elementary?, J. Phys. A: Math. Gen., 37 (2004) 3285-3304.